

Abstract

Stability of Viscous Shock Profiles via Weighted Energy Estimates
Christian Fries

Abstract:

In this talk the use of weighted energy estimates in the proof of stability of viscous shock profiles is shown. A viscous shock profile is a specific traveling wave solution

$$u(x, t) = \phi(x - st)$$

of the hyperbolic conservation law

$$\begin{aligned} u_t + f(u)_x &= u_{xx} \quad x \in \mathbb{R}, u \in \mathbb{R}^n \\ u(\pm\infty) &= u_{\pm} \end{aligned} \tag{1}$$

(f' is assumed to have real eigenvalues, s and u_{\pm} are subject to certain conditions). By stability we mean:

Theorem 1

Let $U_0 \in H^{2,2}$. If $|u_+ - u_-|$ and $\|U_0\|_{H^{2,2}}$ are sufficiently small, then the solution $u(x, t)$ to (1) with data $u(\cdot, 0) = \phi + (U_0)_x$ exists for all times $t > 0$ and has

$$\limsup_{t \rightarrow 0} \sup_x |u(x, t) - \phi(x - st)| = 0$$

In the course of the talk we will present the “integrated equation”, which is the starting point of the energy estimates. Then we will sketch a basic energy estimate for a scalar equation ($n = 1$) using a convexity assumption on f , show how this assumption can be removed by the use of weight functions and finally present the main ingredients for a similar treatment of a system ($n > 1$).